
Species	Interior	Coastal	Species	Interior	Coastal	Species	Interior	Coastal
Hummingbirds			<u>Thrushes</u>			<u>Finches</u>		
Rufous Hummingbird	*R (2)	*C (1,2,3)	American Robin	*C (1,2,3)	*C (1,2,3)	Pine Grosbeak	*U (1,2,3,4)	*U (1,2,3,4)
			Townsend's Solitaire	*C (1,2,3)	*C (1,2,3)	Rosy Finch	*U (1,2,3)	*U (1,2,3)
Kingfishers			Hermit Thrush	*U (1,2,3)	R (1,2,3)	Common Redpoll	*C (1,2,3,4)	*R (1,4)
Belted Kingfisher	*C (1,2,3)	*U (1,2,3,4)	Swainson's Thrush	*U (1,2,3)	*C (1,2,3)	Hoary Redpoll	*C (1,4)	*R (1,4)
<u> </u>	(,	, ,	Gray-cheeked Thrush	*C (1,2,3)	U (1,2,3)	Pine Siskin	*U (1,2,3)	*C (1,2,3)
Woodpeckers			Mountain Bluebird	*R (1,2,3)	+ (3)	Red Crossbill		*R (1,2,3,4)
Northern Flicker	*C (1,2,3)	*U (3)	Northern Wheatear	*U (1,2,3)	R (1,2,3)	White-winged Crossbill	*C (1,2,3,4)	*U (1,2,3,4)
Hairy Woodpecker	*C (1,2,3,4)	*U (1,2,3,4)		- (-,=,-,	(.,=,-,	Brambling	*+ (1,3,4)	+ (3,4)
Downy Woodpecker	*U (1,2,3,4)	*U (1,2,3,4)	Kinglets				(.,5,.)	(5, .)
Black-backed Woodpecker		+ (1,2,3)	Golden-crowned Kinglet	R (2,3)	*U (1,2,3,4)			
Black-backed vvoodpecker Three-toed Woodpecker	*U (1,2,3,4)		Golden-crowned Kinglet Ruby-crowned Kinglet	*C (1,2,3)		If you observe any hinds that	ara not on this sl	hacklist places
		*R (1,2,3,4)	Ruby-crowned Kinglet	··C (1,2,3)	*C (1,2,3)	If you observe any birds that		necklist please
Red-breasted Sapsucker	*+ (2)	+ (1,3,4)	011114/ 11114/ 11			send detailed reports to Par	k Headquarters:	
.			Old World Warblers					
<u>Flycatchers</u>			Arctic Warbler	*R (1,2,3)		Wrangell-St. Elias	National Park and	Prosorvo
Say's Phoebe	*U (1,2,3)	R (1,2,3)					i National Laik and	1 1 6361 46
Alder Flycatcher	*C (1,2,3)	R (1,2,3)	<u>Pipits</u>			PO Box 439		
Hammond's Flycatcher	*U (1,2,3)		American Pipit	*C (1,2,3)	*C (1,2,3)	Copper Center, A	K 99573	
Western Wood Pewee	*U (1,2,3)	R (1,2,3)	•			(907)-822-5234		
Olive-sided Flycatcher	*U (1,2,3)	R (1,2,3)	Waxwings			wrst interpretation	n@nps.gov	
•	, ,	, ,	Bohemian Waxwing	*C (1,2,3)	U (1,2,3)		O 1 - 6	
Larks			Boneman waxwing	C (1,2,3)	0 (1,2,3)			
Horned Lark	*C (1,2,3)	R (1,2,3)	Shrikes			VEV T	SYMBOLS:	•
	J (1,2,5)	(. , = , =)		*!! (! 2 2)	*!! /! 2 2\			
<u>Swallows</u>			Northern Shrike	*U (1,2,3)	*U (1,2,3)	 Positive identification mad 	le in the Park/Pres	erve.
	*C (1 2 2)	C (1.2.2)	6. 1			C Common		
Violet-green Swallow	*C (1,2,3)	C (1,2,3)	<u>Starlings</u>			U Uncommon		
Tree Swallow	*C (1,2,3)	*C (1,2,3)	European Starling	R (1,2,3)	R (1,3,4)			
Bank Swallow	*C (1,2,3)	*U (1,2,3)				R Rare		
Cliff Swallow	*C (1,2,3)	U (1,2,3)	New World Warblers			 + Casual or accidental vag 	grants.	
Barn Swallow	*R (1,2,3)	*C (1,2,3)	Orange-crowned Warbler	*C (1,2,3)	*C (1,2,3)	- Not known to occur w	-	
			Yellow Warbler	*U (1,2,3)	*U (1,2,3)	1400 Kilowii to Occui W	idilii die region	
Jays and Crows			Yellow-rumped Warbler	*C (1,2,3)	*U (1,2,3)			
Gray Jay	*C (1,2,3,4)	*R (1,2,3,4)	Townsend's Warbler	*R (1,2,3)	*U (1,2,3)			
Steller's Jay	+ (3)	*C (1,2,3,4)	Blackpoll Warbler	*U (1,2,3)	*R (1,2,3)	Numbers with parentheses	represent season:	
Common Raven	*C (1,2,3,4)	*C (1,2,3,4)	Northern Waterthrush	*C (1,2,3)	*R (1,2,3)			
Northwestern Crow		*C (1,2,3,4)	Wilson's Warbler	*C (1,2,3)	*R (1,2,3)	I=Spring, 2=Sumr	ner, 3=Fall, 4=VV	/inter
Black-billed Magpie	*C (1,2,3,4)	*C (1,2,3,4)	VVIISOITS VVALDIEI	C (1,2,3)	TK (1,2,3)			
	•	•						
<u>Chickadee</u>			Blackbirds			The two bio-geographic regi	ons considered ar	e:
Black-capped Chickadee	*C (1,2,3,4)	*U (1,2,3,4)	Rusty Blackbird	*U (1,2,3)	*R (1,2,3)			
Chestnut-backed Chickade		*C (1,2,3,4)	Red-winged Blackbird	R (1,2,3)	*R (1,2,3)	Interior = North o	of the Bagley Icefie	ld
Siberian Tit	+ (1,2,3,4)		Brown-headed Cowbird	+ (1,2,3)	+ (1,2,3)	Caracal - Caral	التكميدا بتجامم الممامة	٦
Boreal Chickadee	*C (1,2,3,4)	R (1,2,3,4)	Brown-neaded Cowbiid	. (1,2,3)	. (1,2,3)	Coastal = South o	i trie bagiey icefiel	u
			Sparrows, Juncos, Bunti	ngs				
<u>Nuthatches</u>			Savannah Sparrow	*C (1,2,3)	*C (1,2,3)			
Red-breasted Nuthatch	*+ (1,2,3,4)	*U (3)	Dark-eyed Junco	*C (1,2,3)	*U (1,2,3,4)			
			American Tree Sparrow	*C (1,2,3)	U (1,3)			
<u>Creepers</u>			Chipping Sparrow	R (1,2,3)	+ (1,2,3)	Date		
Brown Creeper	R (1,2,3,4)	*U (1,2,3,4)	Brewer's Sparrow	*+ (1,2)	(1,2,3)	_ ****		
	() / /	(, ,-, ,	White-crowned Sparrow	*C (1,2,3)	*U (1,2,3)	Location		
Wrens			Golden-crowned Sparrow			LOCATION		
Winter Wren		*U (1,2,3,4)		,	*C (1,2,3)	Observer(s)		
vviiitei vvieii		· U (1,2,3, 4)	Fox Sparrow	*C (1,2,3)	*C (1,2,3)	Observer(s)		
D'			Lincoln's Sparrow	*C (1,2,3)	*C (1,2,3)	1		
<u>Dippers</u>		******	Song Sparrow	R (1,2,3)	*C (1,2,3,4)	Remarks		
American Dipper	*U (1,2,3,4)	*C (1,2,3,4)	Lapland Longspur	*C (1,2,3)	U (1,3)			
			Smith's Longspur	R (1,2,3)				
<u>Doves</u>			Snow Bunting	*C (1,2,3)	*U (1,3)			
Mourning Dove	+ (1,2,3)	R (3)						

Wrangell-St. Elias



Bird Checklist





National Park Service U.S. Department of the Interior Wrangell- St. Elias National Park & Preserve

Birds of Wrangell-St. Elias National Park & Preserve

This is a list of 212 bird species found in Wrangell-St. Elias National Park & Preserve. The park contains 13.2 million acres, including both marine shoreline near Yakutat and vast expanses of wild, interior Alaska with boreal spruce forests, muskegs, alder thickets, alpine meadows, icefields, and glacial barrens.

Long summer days and abundant habitats lure long distance migrants through the Copper River Valley each spring. Trumpeter swans, Canada geese, and a myriad of other waterfowl and shorebirds begin arriving in April. By early May, park forests are alive with birdsong as warblers and thrushes establish nesting territories and find mates. By August, many species are already beginning their return to southern wintering grounds.

As days shorten and the frigid winter of the Alaskan interior arrives, only the hardiest 34 species remain. Foraging chickadees, redpolls, and grosbeaks can be seen and heard on all but the coldest of days.

As one of the largest wilderness areas in the world, and with limited road access, this rugged area offers both unique challenges and rich rewards to the casual observer and seasoned birder alike. A journey along either the 60-mile McCarthy Road or 40-mile Nabesna Road will provide for many opportunities to enjoy summer's bird pageantry in spectacular settings.

Stop by any park visitor center to ask about recent sightings and check on road conditions before heading out.

Common Loon	
Common Loon C(1,2) C(1) C(1,3) C(1,3) C(1,3) King Eider	C (1,2,3)
Yellow-billed Loon	C (1,3)
Red-horsated Loon C (1.3) C (1.3.4) Steller's Edder C (1.3) C (1.3.4) Red-horsated Phalarope C (1.3) C (1.3.4) Red-horsated Phalarope C (1.3.3) Red-horsated Phalarope C (1.3.3) Red-horsated Phalarope C (1.3.4) Red-horsated Phalarope Red Horsated	C (1,2,3)
Red-throated Loon	C (1,3)
Black Scoter R (1,2.3) C (1,3.4) Red-necked Phalarope C (1,2.3) Red-necked Final Phalarope C (1,2.3) Red-necked Grebe C (1,2.3) C (1,3.4) C	C (1,3)
White-winged Scoter C (1,23) C (1,3) C	C (1,2,3)
Red-necked Grebe	
Horned Grebe *C (1,2,3) C (1,3,4) Hooded Merganser R (1,2,3,4) *C (1,23,4)	C (1,2,3)
Red-breasted Merganser **Q (1,2,34) **C (1,2,	C (1,3)
Albatrosses	C (I)
Black-floored Albatross	C (1,3)
Black-foored Albatrors	C (1,3,4)
Laysan Albatross R (1,2,3) Birds of Prey Bald Eagle Sald Eagle C (1,2,3,4) C (1,2,3,4) Dunlin + (1,2,3)	C (1,3)
Bald Eagle C (1,2,3) C (1,2,3,4) Dunin + (1,1,3)	C (I)
Northern Fulmar	C (1,3)
Northern Fulmar	U (1,2,3)
Pink-footed Shearwater	U (1,3)
Soory Shearwater	U (1,3)
Soury start alled Shearwater	C (1,2,3)
Short-tailed Shearwater C (2)	C (1,3)
Storm-petrels	
Falcons	
Storm-petrel + (3) C (1,2,3) Falcons - (2,1,2) Falcons - (2,1,2,3) Falcons - (2,1,	C (2,3)
Fork-tailed Storm-petrel	C (1,3)
Leach's Storm-petrel	R (1,2,3)
Peregrine Falcon	R (1,2,3,4)
Cormorants	C (1,2,3,4)
	C (1,2,3, 1)
Pelagic Cormorant	R (1,2,3,4)
Red-faced Cormorant	R (1,2,3,4)
Spruce Grouse C (1,2,3,4) Spruce Grouse C (1,2,3,4) R (1,2,3,4) Spruce Grouse R (1	C (1,2,3,4)
Ruffed Grouse R (1,2,3,4) Bonaparte's Gull *U (1,2,3) Sabrie's Gull *U (1,2,3) Sabr	
Willow Ptarmigan *C (1,2,3,4) *U (1,2,3,4) Sabine's Gull + (2)	C (1,2,3)
Rock Ptarmigan *C (1,2,3,4) Arctic Tern *U (1,2,3) Arctic Tern	C (1,2,3)
White-tailed Ptarmigan Sharp-Tailed Grouse Scott Grouse Sco	U ((1,3)
Sharp-Tailed Grouse *C (1,2,3,4)	C (1,2,3)
	U (1,2,3)
Trumpeter Swan	
Brant	C (1,2,3,4)
Emperor Goose	R (1,2,3,4)
	C (1,2,3,4)
Greater White-fronted Goose	U (1,2,3)
	C (1,2,3)
Gadwall R (1,2,3) C (1,3) Plovers — Crested Auklet + (3) Northern Pintail *C (1,2,3) *C (1,2,3) — Semi-palmated Plover *C (1,2,3) *C (1,2,3) — Cassin's Auklet Green-winged Teal *C (1,2,3) *C (1,2,3) — Killdeer R (1,2,3) R (1,2,3) — Marbled Murrelet Rive winged Teal *I (1,2) *P (1,2,3) — Lesser Golden Plover *C (1,2,3) — Kittlitz's Murrelet Rive winged Teal *I (1,2) *P (1,2,3) — Kittlitz's Murrelet Green-winged Teal *I (1,2) *P (1,2,3) — Kittlitz's Murrelet Rive winged Teal *I (1,2) *P (1,2,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) *C (1,2,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) *C (1,2,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) *C (1,2,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) *C (1,2,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) *C (1,2,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) *C (1,2,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) *C (1,2,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) *C (1,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) *C (1,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) *C (1,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) *C (1,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) *C (1,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) *C (1,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) *C (1,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) *C (1,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) *C (1,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) *C (1,3) — Kittlitz's Murrelet Semi-palmated Plover *C (1,2,3) — Kittlitz's Murrelet	R (1,2,3,4)
	R (4)
	R (1,2,3)
Green-winged Teal *C (1,2,3) *C (1,2,3)	C (1,2,3,4)
	C (1,2,3)
— Committee of the comm	R (1,2,3,4)
Northern Shoveler *C (1,2,3) *C (1,2,3) = Black-bellied Flover * (1,2,3) = C (1,3) = Parakeet Auklet	R (1,2,3)
American Wigeon	(-,=,-)
Eurasian Wigeon *R (I) R (I) Oystercatchers	
Canvasback *U (1,2,3) C (1,2,3) American Black Oystercatcher *C (1,2,3) Owls	
King-necked Duck	C (1 2 2 4)
	C (1,2,3,4)
	U (4)
Redbead *I (1.3) *R (1.23) \\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	C (4)
Common Goldeneve *C (1.2.3) *C (1.2.3)	R (1,2,3,4)
Barrow's Goldenave *C (1.2.3) *C (1.2.3.4)	C (1,2,3)
Bufflehead	U (1,2,3,4)